

INTRODUCTION

1

This report presents the results of an archeological survey of ca. 991 acres (401 hectares) and the National Register testing of prehistoric site 41BX1180 and historic military training sites 41BX1163 and 41BX1189 by Prewitt and Associates, Inc. Sites 41BX1180, 41BX1163, and 41BX1189 were recorded in 1995–1996 by Kibler et al. (1998) and were assessed as potentially eligible for listing in the National Register of Historic Places. The current survey covers the remaining acreage not covered during the prior phase of work. This work was conducted from 6 January 1997 to 5 March 1997 at Camp Stanley Storage Activity under contract with the U.S. Army Corps of Engineers, Fort Worth District. Camp Stanley is a U.S. Army installation encompassing ca. 4,004 acres in northern Bexar County, Texas (Figure 1); it is divided into an Inner Cantonment (ca. 1,780 acres) and Outer Cantonment (ca. 2,224 acres). The current survey covered acreage in the Outer Cantonment; six archeological sites, consisting of one prehistoric component, one historic premilitary component, and five historic military components, were recorded. Test excavations were performed at 41BX1180, 41BX1163, and 41BX1189 to identify the sites' components and to evaluate their significance in terms of the criteria specified by the U.S. Department of the Interior, National Park Service, National Register Division (1982:1). These investigations were authorized by and conducted in accordance with Section 110 (16 U.S.C. 470h-2(a)(2)) of the National Historic Preservation Act of 1966 as amended through 1992 (P.L. 102-575) and Army Regulation (AR) 200-4 (*Cultural Resources Management*).

The results of these investigations are presented in the following manner. The remainder of Chapter 1 provides environmental background on the geology, Quaternary geomorphology, climate, and flora and fauna of the area. Chapter 2 summarizes the cultural chronologies of the area, including the prehistoric and historic periods. Chapter 3 details the methods of investigations for the pedestrian survey and archeological testing. The results of the investigations are pre-

sented in Chapter 4 with individual site descriptions and summaries as well as analytic results from data collected during test excavations. Assessments and recommendations for the cultural resources are given in Chapter 5.

ENVIRONMENTAL BACKGROUND

Geology

Camp Stanley is located in south-central Texas along the Balcones fault zone, which marks the southern and eastern margins of the Edwards Plateau separating it from the Blackland Prairie and Gulf Coastal Plain to the southeast (Arbingast et al. 1973:6; Brown et al. 1974). The Balcones fault zone is a band of northeast-southwest-trending faults that were active during the Miocene. The fault blocks throughout this zone are generally downthrown to the southeast, a result of the uplift of the Edwards Plateau with respect to the Blackland Prairie and the Gulf Coastal Plain. Camp Stanley is underlain by the Lower Cretaceous Glen Rose Formation (Brown et al. 1974). The Glen Rose Formation consists of alternating beds of limestones, dolomites, and marls indicative of shallow sub-tidal and tidal-flat environments. These alternating resistant and recessive beds give the Glen Rose Formation its characteristic stairstepped or benched topography. The Glen Rose is divided into upper and lower units. The lower Glen Rose is exposed across Camp Stanley in the Salado Creek valley, its tributaries, and smaller drainages (Collins 1994). It is 200–270 ft thick and consists of massive limestones and dolomitic limestones containing some rudistid reefs and mounds. The top of the lower Glen Rose unit is capped by a series of 1–3-ft-thick *Corbula* beds consisting of abundant steinkerns of the bivalve *Corbula harveyi* (Collins 1994). The upper Glen Rose is present on the hilltops, ridges, slopes, and intervening saddles across Camp Stanley and consists of thinner, more-dolomitic, and less-fossiliferous beds (Collins 1994). It has an overall thickness of 400 ft.




Figure 1. Project location map.

Quaternary Geomorphology

The rugged and highly dissected landscape of Camp Stanley is the result of millions of years of erosion. The uplift of the Edwards Plateau during the late Tertiary created knickpoints which streams draining the plateau have been wearing down since the Miocene. Much of Camp Stanley is drained by Salado Creek, an intermittent upland tributary of the San Antonio River. The extreme southwestern corner of the installation is drained by an unnamed ephemeral tributary of Leon Creek, while the northeastern corner is drained by ephemeral tributaries of Cibolo Creek. Faulting has controlled or strongly influenced the development of these drainages (particularly Salado Creek) by directing groundwater to seeps and springs, exposing formations of varying resistances, and controlling gradients.

The low-order upland drainages are usually deep and narrow, and are often scoured to bedrock and devoid of thick alluvial deposits. Relatively thick alluvial deposits (up to 1–2 m thick) were observed only in the Salado Creek valley in the form of a broad floodplain. These deposits consist of thick gravelly basal units capped by thin mantles of dark loamy and clayey alluvium. Much of the Salado Creek valley has been modified and channelized, disturbing and destroying the stratigraphy of the alluvial deposits. Downstream, in Camp Bullis, a large earthen dam was constructed by the Soil Conservation Service on Salado Creek to inhibit rapid run-off and to facilitate the infiltration of water into the Edwards aquifer. Undoubtedly the presence of the dam has affected sedimentation upstream in Camp Stanley. In the past, the steep gradient of upper Salado Creek (9.5 m/km), coupled with heavy rainfalls and impervious ground cover throughout the watershed, resulted in intensive flash floods and scouring of the channel bed and margins. It is probable that most of the upper Salado Creek valley was periodically scoured out, making the presence of any alluvial deposits temporary features within the valley. How often this may have occurred is not known, but since flashy discharges are now controlled to some extent, it suggests that much of the sediment upstream from the dam may have been deposited since its construction in the 1970s.

An earlier archeological survey along Salado Creek in Camp Bullis also noted the presence of basal gravels (high-energy environments) and suggested that most of the valley had little potential for yielding intact archeological sites (Quigg 1988:26). This notion was further supported by the absence of prehistoric cultural resources in primary context throughout the Salado Creek valley even though the area was subjected to an

intensive shovel testing program during the survey.

Uplands, ridges, slopes, and intervening saddles characterize much of the Camp Stanley landscape and therefore represent an important part of the evolution of this landscape and potential archeological record. Kibler (Kibler and Gardner 1997) presents a simple model of landscape evolution for the Camp Stanley area, based on investigations at neighboring Camp Bullis, that primarily focuses on the development of upland slopes and processes of sediment transport from upland environments to the fluvial systems in the valley bottoms. The model views sediments and clasts as being delivered to the valley bottoms from the uplands and slopes by sheetwash, rillwash, fluvial, and mass-wasting processes through a system of discontinuous and segmented bedrock-entrenched gullies. Other colluvial processes and depositional environments commonly associated with the development of slopes, such as debris flows and the formation of fans, are extremely rare.

Upland or upslope sediments are delivered by sheetwash and rillwash to the gullies, where the flow becomes channelized and restricted. The gullies tend to be widely spaced, creating broad, smooth slopes characteristic of a system of low drainage density (Young 1972:74). The alternating resistant and pervious beds of the Glen Rose Formation, as well as the gradient of the main slope, play a major role in the development of these gullies, influencing and controlling drainage, channel (gully) morphology, gradient, and incision. Farther downslope, the walls and bottom of the channel merge at a location known as the intersection point (Waters 1992:145). Intersection points usually occurred where the drainage intersects an exposed bed of highly resistant bedrock, hindering entrenchment, and where the gradient of the main slope decreases. Sediments accumulate at intersection points to form small gully-mouth fans, sheetwash and rillwash become the dominant modes of sediment transport, and the overall gully channel becomes wider and shallower until the channel once again becomes entrenched due to a bedrock or gradient change downslope. Channel gradient changes tend to be extremely sharp, localized, and marked by <1–3-m-high pour-offs, which denote changes in the resistance of the exposed bedrock. Entrenchment is usually through headward erosion and the undercutting and mass wasting of more-resistant limestone caps at the pour-offs.

Depositional environments on the uplands and valley slopes are rare. These surfaces are most often nonaggrading or deflationary with very thin soil mantles, primarily the result of in situ weathering or deposition through sheetwash and rillwash. The few

depositional environments existing on the slopes are the result of sediments filling local topographic depressions, accumulating behind barriers (e.g., vegetation), or accumulating where gradients sharply decline (e.g., intersection points). These deposits tend to be thin (rarely more than 40 cm thick) discontinuous patches of gravelly loams.

Soils throughout Camp Stanley belong to the Tarrant-Brackett association (Taylor et al. 1966:2). A variety of soil series and types are included in this association; however, the thin, gravelly, calcareous Tarrant and Brackett soils are predominant, mantling the hilltops and slopes. Reddish brown and gravelly Crawford and Bexar soils occupy the lower valley slopes and intervening saddles. Dark-colored, calcareous, and clayey Krum soils occupy the toeslopes of the larger valleys and low-order upland drainages, while the dark-colored, clayey Trinity and Frio soils occupy the floodplain of Salado Creek.

Climate

The climate of south-central Texas can be classified as modified humid subtropical with Gulf-influenced hot summers and continental-influenced mild winters. Mean daily maximum and minimum temperatures for January are 62°F and 39°F, respectively. For July, mean daily maximum and minimum temperatures are 95°F and 74°F, respectively. The mean annual precipitation for the region is 81.2 cm (32 inches), but varies west to east across the region from 66 to 106 cm. Precipitation falls throughout the year but is generally in the form of late spring and early fall thunderstorms (Natural Fibers Information Center 1987:12, 49–50). The movement of warm, moist Gulf air up onto the edge of the Edwards Plateau during warmer months often triggers short-lived but heavy cloudbursts, making the region one of the most flood-prone in the United States (Patton and Baker 1977).

Flora and Fauna

Camp Stanley is located along the southern margin of the Balconian biotic province (Blair 1950). However,

the floral and faunal assemblages tend to represent a mix of three biotic provinces—the Balconian, Texan, and Tamaulipan. The area also represents the geographical limit for the distribution of many faunal species (Blair 1950:113–114).

Early historic accounts of the floral community describe upland grassland savannahs with dense woodlands along stream valleys (Bonnell 1840:94; Roemer 1935:118). However, development, overgrazing, and suppression of natural range fires throughout the region in the nineteenth and twentieth centuries have severely altered vegetation patterns. Today, upland areas consist of dense scrub forests of juniper (*Juniperus ashei*), Texas oak (*Quercus texana*), and live oak (*Q. virginiana*) with scattered Texas redbud (*Cercis texensis*), prickly pear (*Opuntia* sp.), beargrass (*Nolina texana*), and sotol (*Dasilyrion* sp.) as understory vegetation. Dense patches of sotol and beargrass are present in upland and slope areas that have been subjected to recent (within the last 2–3 years) prescribed burnings. Along the stream valleys, a diverse number of species are present, including live oak, hackberry (*Celtis laevigata*), cedar elm (*Ulmus crassifolia*), and pecan (*Carya illinoensis*). These riparian zones may be interspersed with open grassy areas of little bluestem (*Schizachyrium scoparium*) and other tall grasses (Blair 1950:112–113).

A diverse number of faunal species occupy the area due to the intersection of three biotic provinces and because of the diverse number of habitats and microenvironments along the edge of the Edwards Plateau (Lundelius 1967:309–310). Present-day species include white-tailed deer (*Odocoileus virginianus*), coyote (*Canis latrans*), gray fox (*Urocyon cinereoargenteus*), bobcat (*Lynx rufus*), raccoon (*Procyon lotor*), striped skunk (*Mephitis mephitis*), fox squirrel (*Sciurus niger*), eastern cottontail rabbit (*Sylvilagus floridanus*), turkey (*Meleagris gallopavo*), and 49 other mammals, 36 species of snakes, and 16 lizard species (Blair 1950:113–115). Prehistorically, important economic species of the area included bison (*Bison bison*), pronghorn antelope (*Antilocapra americana*), and black bear (*Ursus americanus*) (Gerstle et al. 1978:29).